




THEOREM OF THE DAY

The Second Isomorphism Theorem Suppose H is a subgroup of group G and K is a normal subgroup of G . Then HK is a group having K as a normal subgroup, $H \cap K$ is a normal subgroup of H , and there is an isomorphism from $H/(H \cap K)$ to HK/K defined by $h(H \cap K) \mapsto hK$.

Cayley graph of Frobenius group F_{20}



KEY

$$a := (12345)$$

$$b := (1254)$$

Vertices (group elements)

$$(m,n) := b^m a^n = (1254)^m (12345)^n$$

$$\text{E.g. } (1,2) = b^1 a^2$$

$$= (1254)(12345)^2$$

$$= (1254)(13524)$$

$$= (1435)$$

Edges (left multiplication)

→ by a

— by b


(anticlockwise cycle)

To illustrate we take G to be Sym_5 , the group of $5!$ permutations of $\{1, 2, 3, 4, 5\}$. The Frobenius group F_{20} may be defined as a subgroup H of G generated by a 5-cycle, a , and a 4-cycle, b , satisfying $(ab)^4 = a(ab)(ba)^{-1} = 1$. We take K to be Alt_5 , the subset of $5!/2$ even permutations: identity, 5-cycles, and products of two 2-cycles. This is normal in G (i.e. $g^{-1}Kg = K$ for all g) because conjugation, $g^{-1}xg$, is a 1-1 mapping which preserves cycle structure; similarly, $H \cap K$, the subset of even permutations in H , is a normal subgroup of H .

Now we can uncover the behaviour of the normal subgroup of even permutations of F_{20} . The target quotient, HK/K , is $\text{Sym}_5/\text{Alt}_5 \cong C_2$ because (1) any coset of K , say, hK , is either all of K (if h is even) or all odd permutations in Sym_5 (if h is odd), and (2) multiplication of cosets mirrors addition modulo 2: $hK \cdot h'K = hh'K$ switches coset if and only if h and h' have different parity. So cosets of $H \cap K$ must behave in exactly the same way in H . And we can see this in the Cayley graph: $H \cap K$ is the 'identity' coset consisting of all vertices (m,n) , m even; there is one other coset, $b(H \cap K)$, and left multiplication by b cycles between the two.

This theorem, due in its most general form to Emmy Noether in 1927, is an easy corollary of the first isomorphism theorem. Thus, if $f : H \rightarrow HK/K$ is the surjective homomorphism $h \mapsto hK$ then $H/\ker f \cong \text{im} f$ and $\ker f = H \cap K$.

It is sometimes called the 'parallelogram rule' in reference to the diagram on the right.



Web link: people.reed.edu/~jerry/332/09isom.pdf

Further reading: *Classic Algebra* by P.M. Cohn, John Wiley & Sons, 2000, Chapter 9.

