





THEOREM OF THE DAY

Green's Theorem Let C be a closed, anticlockwise-oriented curve in the xy -plane enclosing a region D . Let $F(x, y) = (P(x, y), Q(x, y))$ be a 2-valued function having continuous partial derivatives on C and inside D . Then

$$\int_C F ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$



Suppose we think of F as a force field acting in the plane. Then the line integral $\int_C F ds$ may be thought of as measuring the total work done by F acting on a particle as it follows the curve C ; the integral is often written in the form $\int_C P dx + Q dy$, making explicit the action of the x and y components of the force as the particle moves through a small increment in the x and y directions. We refer to this work done by F as the 'circulation of F around C '. Green's Theorem asserts that circulation around C is the accumulation of 'microscopic circulations' around points in D : see the illustration on the left; these microscopic circulations are measured as the component perpendicular to the plane of the **curl** of F ; this component is calculated as: $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$.



As an illustration we take $P(x, y) = (x - y)^2$ and $Q(x, y) = x + y$. These are plotted as surfaces in 3D on the right, and a half-unit circle, closed by adjoining a segment of the x -axis, is chosen as the curve C . We find $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 1 + 2(x - y)$ and Green's Theorem yields the value of $\int_C F ds$ by double integration as

$$\iint_D 1 + 2(x - y) dA = \int_{-1}^1 \int_0^{\sqrt{1-x^2}} 1 + 2(x - y) dy dx = \int_{-1}^1 \left(\sqrt{1-x^2} + 2x\sqrt{1-x^2} - 1 + x^2 \right) dx = \left[\frac{1}{2}x\sqrt{1-x^2} + \frac{1}{2}\sin^{-1}x - \frac{2}{3}(1-x^2)^{3/2} - x + \frac{x^3}{3} \right]_{-1}^1 = \frac{\pi}{4} - \frac{4}{3}.$$

Evaluating the line integral directly requires piecewise parameterisation of the curve C : half-circle $c_1(t) = (-\cos t, \sin t)$, $0 \leq t \leq \pi/2$ and base $c_2(t) = (t, 0)$, $-1 \leq t \leq 1$:

$$\begin{aligned} \int_C F ds &= \int_0^{\pi/2} F(c_1(t)) \cdot c_1'(t) dt + \int_{-1}^1 F(c_2(t)) \cdot c_2'(t) dt = \int_0^{\pi/2} F(\cos t, \sin t) \cdot (-\sin t, \cos t) dt + \int_{-1}^1 F(t, 0) \cdot (1, 0) dt \\ &= \int_0^{\pi/2} (\cos t - \sin t)^2(-\sin t) + (\cos t + \sin t) \cos t dt + \int_{-1}^1 t^2 dt = -2 + \frac{\pi}{4} + \frac{2}{3} = \frac{\pi}{4} - \frac{4}{3}, \text{ as expected.} \end{aligned}$$

George Green published this theorem, a powerful generalisation of the Fundamental Theorem of the Calculus, in 1828.

Web link: mathinsight.org/greens_theorem_idea (on which the above description and examples are based).

Further reading: *Inside Interesting Integrals* by Paul J. Nahin, Springer, 2015, Chapter 8.

Created by Robin Whitty for www.theoremoftheday.org

