




THEOREM OF THE DAY

Euler's Continued Fraction Correspondence Let $(a_i)_{i \geq 0}$ be an infinite sequence of nonzero real or complex numbers. Let f_n denote the n -th partial sum of the sequence: $f_n = \sum_{i=0}^n a_i$. Then f_n is also the n -th convergent of the continued fraction

$$a_0 + \frac{a_1}{1} + \frac{-a_2/a_1}{1 + a_2/a_1} + \dots + \frac{-a_n/a_{n-1}}{1 + a_n/a_{n-1}} + \dots$$



An infinite summation due to the 15th century Indian scholar Kerala Gargya Nilakantha:

$$\frac{\tau - 6}{4} = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k(2k+1)(2k+2)}$$

$$= \frac{1}{1.3.4} - \frac{1}{2.5.6} + \frac{1}{3.7.8} - \frac{1}{4.9.10} + \dots$$

If we apply Euler's correspondence to Nilakantha's series with $a_i = (-1)^{k-1}/k(2k+1)(2k+2)$ then we get $\frac{\tau - 6}{4} = \frac{1}{12} + \frac{1^2 \cdot 3^2 \cdot 4^2}{12 \cdot 2^2} + \frac{2^2 \cdot 5^2 \cdot 6^2}{12 \cdot 3^2} + \frac{3^2 \cdot 7^2 \cdot 8^2}{12 \cdot 4^2} + \dots$, giving $\tau = 6 + \frac{4}{12} + \frac{1^2 \cdot 3^2 \cdot 2^2 \cdot 2^2}{12 \cdot 2^2} + \frac{2^2 \cdot 5^2 \cdot 2^2 \cdot 3^2}{12 \cdot 3^2} + \frac{3^2 \cdot 7^2 \cdot 2^2 \cdot 4^2}{12 \cdot 4^2} + \dots = 6 + \frac{2^2}{12} + \frac{6^2}{12} + \frac{10^2}{12} + \dots$, whose convergents are explored in the above pie chart.

Leonhard Euler discovered this correspondence in 1748. The above application to τ (in a π version) was given by Douglas Bowman as an alternative derivation of a continued fraction published by Jerome Lange in 1999.

Web link: people.math.binghamton.edu/dikran/478/Ch7.pdf.

Further reading: *Handbook of Continued Fractions for Special Functions* by Annie Cuyt et al, Springer, 2008.

